THE JOURNAL OF CHEMICAL THERMODYNAMICS

Volume 16, 1984

EDITORS

M. L. McGlashan
Department of Chemistry
University College London
20 Gordon Street
London WC1H 0AJ
U.K.

P. A. G. O'Hare
Argonne National Laboratory
Chemical Technology Division
Argonne
Illinois 60439
U.S.A.

Assistant Editor: **M. B. Ewing** (University College London, U.K.)

ADVISORY BOARD

Chairman: G. M. Schneider (F.R.G.)

D. Ambrose (U.K.)
G. C. Benson (Canada)
M. Colomina (Spain)
C. Jolicoeur (Canada)
O. J. Kleppa (U.S.A.)
M. Månsson (Sweden)
V. A. Medvedev (U.S.S.R.)
F. L. Oetting (U.S.A.)
J. B. Ott (U.S.A.)
F. Pavese (Italy)
G. Pilcher (U.K.)
J. Stecki (Poland)
H. Suga (Japan)
F. L. Swinton (U.K.)
A. G. Williamson (N.Z.)

ACADEMIC PRESS
London Orlando San Diego
San Francisco New York Toronto
Montreal Sydney Tokyo São Paulo
A Subsidiary of Harcourt Brace Jovanovich, Publishers

Copyright © 1984 Academic Press Inc. (London) Limited

ALL RIGHTS RESERVED

No part of this volume may be reproduced in any form, by photostat, microfilm, or any other means, without written permission from the publishers.

Printed in Great Britain

CONTENTS OF VOLUME 16

Number 1, January 1984

Editorial Announcement i

GOVINDAN, A. P.; VARMA, Y. B. G.; ANANTH, M. S. Isothermal (vapor + liquid) equilibria of four binary mixtures 1

LAÍNEZ, A.; ESCUDERO, I.; DÍAZ PEÑA, M. Excess functions of (1-bromobutane + ethylbenzene) at various temperatures 7

SACCHETTO, G.; KODEJŠ, Z. Vapour pressures of very concentrated electrolyte solutions. III. Measurements on $[(1-x)(CH_3)_2SO + x\{(1-y)NH_4NO_3 + yLiNO_3\}]$ by a dew-point apparatus 15

RUIZ, B.; OTÍN, S.; GUTIÉRREZ LOSA, C. Excess molar enthalpies at 298.15 K of chlorobenzene +, (chloromethyl)benzene +, and (2-chloroethyl)-benzene + an ether 25

TAMURA, K.; MURAKAMI, S. Speeds of sound, isentropic and isothermal compressibilities, and isochoric heat capacities of $\{xc\text{-}C_6H_{12}+(1-x)C_6H_6\}$ from 293.15 to 303.15 K 33

OLOFSSON, G. Temperature dependence of the heat-capacity change for the dissociation of acetic acid and of propionic acid in water. 39

O'HARE, P. A. G.; HUBBARD, W. N.; JOHNSON, G. K.; FLOTOW, H. E. Calorimetric measurements of the low-temperature heat capacity, standard molar enthalpy of formation at 298.15 K, and high-temperature molar enthalpy increments relative to 298.15 K of tungsten disulfide (WS_2), and the thermodynamic properties to 1500 K 45

MILTENBURG, J. C. VAN; BOUWSTRA, J. A. Thermodynamic properties of *trans*-azobenzene and *trans*-stilbene 61

GROLIER, J.-P. E.; INGLESE, A.; WILHELM, E. Excess molar heat capacities of (1,4-dioxane + an *n*-alkane): an unusual composition dependence . . 67

BRAVO, R.; PINTOS, M.; BALUJA, M. C.; PAZ ANDRADE, M. I.; ROUX-DESGRANGES, G.; GROLIER, J.-P. E. Excess volumes and excess heat capacities of some mixtures: (an isomer of hexanol + an *n*-alkane) at 298.15 K 73

FRANZOSINI, P.; WESTRUM, E. F., JR. Thermophysics of metal alkanoates. III. Heat capacities and thermodynamic properties of lithium and potassium propanoates 81

Skudlarski, K.; Kapała, J. Thermodynamics of the sublimation of thallium halides investigated by the mass-spectrometric method 91

Number 2, February 1984

Lavut, E. G.; Timofeyev, B. I.; Yuldasheva, V. M. Enthalpy of formation of iron(III) chloride 101

Navarro, J. M.; Pintos, M.; Bravo, R.; Paz Andrade, M. I. Excess enthalpies of (propyl ethanoate or ethyl propanoate + an *n*-alkane) at 298.15 K 105

Jain, D. V. S.; Sidhu, R. S. Excess molar Gibbs free energies of methylcyclohexane + 1,4-dioxane and + tetrahydrofuran and of ethylbenzene + 1,4-dioxane and + tetrahydrofuran at 303.15 and 323.15 K 111

Wieczorek, S. A. Determination of the excess molar volumes of (butan-2-ol + *n*-heptane) between 293.15 and 308.15 K by use of a continuous dilution dilatometer 115

Tamura, K.; Ohomuro, K.; Murakami, S. Speeds of sound, densities, and isentropic compressibilities of $\{xC_7H_{16}+(1-x)C_6H_6\}$ at 293.15 and 303.15 K 121

Franzosini, P.; Westrum, E. F., Jr. Thermophysics of metal alkanoates. IV. Heat capacities and thermodynamic properties of potassium 2-methylpropanoate 127

Awwad, A. M.; Pethrick, R. A. Isentropic compressibilities of hydrocarbons and their mixtures. Mixtures of linear and branched-chain alkanes . . 131

Ribeiro da Silva, M. A. V.; Ribeiro da Silva, M. D. M. C.; Carvalho, A. P. S. M. C.; Akello, M. J.; Pilcher, G. Standard enthalpies of formation of bis(pentane-2,4-dionato)Cu(II) and of four bis(methyl-substituted heptane-3,5-dionato)Cu(II) complexes: the mean (Cu–O) bond-dissociation enthalpies 137

Tamura, K.; Kumaran, M. K.; Halpin, C. J.; Benson, G. C. Ultrasonic speeds and isentropic compressibilities of each of the mixtures: (2-ethylbutan-1-ol + an isomer of hexane) at 298.15 K 145

Chylinski, K.; Stryjek, R. Excess molar volumes of *cis*-bicyclo[4.4.0]decane + tetrachloromethane, + benzene, + *n*-octane, + *n*-undecane, + *n*-hexadecane, and + *trans*-bicyclo[4.4.0]decane; and of *trans*-bicyclo[4.4.0]decane + tetrachloromethane and + benzene at 298.15 K . . 153

Cibulka, I.; Nguyen, V. D.; Holub, R. Excess molar volumes of (an alkanol + acetonitrile) at 298.15 and 308.15 K 159

STAVELEY, L. A. K.; WEIR, R. D. The heat capacity of potassium metaperiodate, KIO_4, from 8.0 to 309 K 165

SAKURAI, M.; NAKAGAWA, T. Densities of dilute solutions of water in *n*-alkanols at 278.15, 288.15, 298.15, 308.15, and 318.15 K. Partial molar volumes of water in *n*-alkanols 171

KUMARAN, M. K.; BENSON, G. C. Excess enthalpies of decan-1-ol + *n*-pentane, + *n*-hexane, + *n*-heptane, + *n*-octane, + *n*-nonane, + *n*-decane, and + *n*-undecane at 298.15 K 175

KUMARAN, M. K.; BENSON, G. C. Excess molar volumes of (benzene + cyclohexane) at 323.15 K 183

FRANCESCONI, A. Z. Values of (p, V_m, T) for methane in the range 15 to 270 MPa and 323 to 723 K 189

NOTES

LAVUT, E. G.; TIMOFEYEV, B. I.; YULDASHEVA, V. M. Calorimetric determination of $(\partial U_m/\partial p)_{T=298.15\,K}$ for Cl_2 and consequent correction to $\Delta_f H^\circ_m(NbCl_5, s, 298.15\ K)$ 195

BERNAL, P.; VAN HOOK, W. A. Isentropic compressibility of benzene-h_6 and benzene-d_6. 197

Number 3, March 1984

NASIR, P.; SIVARAMAN, A.; KOBAYASHI, R. The molar volume and fugacity of liquid dibenzofuran at high temperatures and high pressures . . . 199

BALDUCCI, G.; GIGLI, G.; GOLDENBERG, M. M. F.; GUIDO, M. Vaporization of $CoTiO_3$ 207

WÓYCICKI, W. Excess enthalpies of binary mixtures containing unsaturated hydrocarbons 219

ZEGERS, H. C.; SOMSEN, G. Partial molar volumes and heat capacities in (dimethylformamide + an *n*-alkanol) 225

TRESZCZANOWICZ, A. J.; BENSON, G. C. Excess volumes for (an *n*-alkanol + an *n*-alkane). V. (Dodecan-1-ol + *n*-heptane) 237

KAMBAYASHI, S.; KATO, E. A thermodynamic study of (magnesium oxide + silicon dioxide) by mass spectrometry at 1973 K 241

CHRISTENSEN, J. J.; CHRISTENSEN, S. P.; SCHOFIELD, R. S.; FAUX, P. W.; HARDING, P. R.; IZATT, R. M. The excess enthalpies of (carbon dioxide + pyridine) at 308.15, 358.15, and 413.15 K from 7.50 to 12.50 MPa . . 249

LIBUŚ, Z.; ZAK, E.; SADOWSKA, T. Osmotic and activity coefficients of aqueous rare-earth ethylsulphates 257

POZHARSKAYA, G. I.; KASAPOVA, N. L.; SKRIPOV, V. P.; KOLPAKOV, YU. D. The spinodal approximation by the method of scattering of light in (*n*-hexane + *n*-tetradecafluorohexane) 267

CROUCH-BAKER, S.; DAVIES, P. K.; DICKENS, P. G. Standard molar enthalpies of formation of each of a series of molybdates: $K_2Mo_nO_{3n+1}$ with $n = 1$, 2, 3, 4, and 8, by solution calorimetry 273

ATAKE, T.; FUJIWARA, T.; CHIHARA, H. A thermal anomaly in solid hexamethylbenzene-d_{18} below 20 K 281

RAU, H. The chlorination equilibrium of germanium oxides 287

JOHNSON, G. K. Enthalpy increments relative to 298.15 K of RbCl(cr) and RbCl(l) 295

NOTE

CROUCH-BAKER, S.; DICKENS, P. G. Standard molar enthalpy of formation of α-$MoO_3 \cdot H_2O$ by solution calorimetry 301

Number 4, April 1984

ROY, R. N.; GIBBONS, J. J.; WILLIAMS, R.; GODWIN, L.; BAKER, G.; SIMONSON, J. M.; PITZER, K. S. The thermodynamics of aqueous carbonate solutions. II. Mixtures of potassium carbonate, bicarbonate, and chloride . . 303

NILSSON, S.-O.; WADSÖ, I. A flow-microcalorimetric vessel for solution of small quantities of easily or slightly soluble liquids. Solution of benzene in water at 298.15 K 317

STARZEWSKI, P.; WADSÖ, I.; ZIELENKIEWICZ, W. Enthalpies of vaporization of some *N*-alkylamides at 298.15 K 331

MURRAY, W. T.; O'HARE, P. A. G. Thermochemistry of inorganic sulfur compounds. II. Standard enthalpy of formation of germanium disulfide . 335

BUSEY, R. H.; HOLMES, H. F.; MESMER, R. E. The enthalpy of dilution of aqueous sodium chloride to 673 K using a new heat-flow and liquid-flow microcalorimeter. Excess thermodynamic properties and their pressure coefficients. 343

BICKERTON, J.; PILCHER, G.; AL-TAKHIN, G. Enthalpies of combustion of the three aminopyridines and the three cyanopyridines 373

COLOMINA, M.; JIMENEZ, P.; ROUX, M. V.; TURRION, C. Thermochemistry of *p*-ethylbenzoic acid 379

RODRIGUEZ-NUÑEZ, E.; PAZ-ANDRADE, M. I.; NAVARRO, J. M. Thermodynamic properties of (a xylene + an *n*-alkanol). I. Excess molar enthalpies at 298.15 K for 1-hexanol, 1-heptanol, and 1-octanol . . 385

EASTEAL, A. J.; WOOLF, L. A. Measurement of (p, V, x) for (ethanol + trichloromethane) at 298.15 K 391

NOTES

HOLDER, G. D.; KAMATH, V. A. Hydrates of (methane + *cis*-2-butene) and (methane + *trans*-2-butene) 399

PILCHER, G.; BICKERTON, J.; RIBEIRO DA SILVA, M. M.D.C.C. Certification of B.D.H. Thermochemical Standard Benzoic Acid 401

Number 5, May 1984

BOUWSTRA, J. A.; OONK, H. A. J.; BLOK, J. G.; KRUIF, C. G. DE. Properties of mixed crystalline organic material prepared by zone levelling. III. Vapour pressures of (*trans*-azobenzene + *trans*-stilbene) 403

INABA, H.; MIMA, T.; NAITO, K. Measurement of enthalpies of formation of niobium oxides at 920 K in a Tian-Calvet-type calorimeter . . . 411

WEINTRAUB, R.; APELBLAT, A.; TAMIR, A. Enthalpy of solution of rubidium nitrate in water 419

CARSON, A. S.; SPENCER, J. A. The enthalpies of formation of bis-benzyl mercury and bis-phenylethynyl mercury and some associated bond-dissociation enthalpies 423

CARSON, A. S. The molar enthalpy of dehydration of $Cr_2(O_2CCH_3)_4 \cdot 2H_2O$(cr) and the molar enthalpies of sublimation of $Cr_2(O_2CCH_3)_4$(cr), $Mo_2(O_2CCH_3)_4$(cr), and $Mo(O_2CCF_3)_4$(cr) 427

AMITIN, E. B.; MINENKOV, YU. F.; NABUTOVSKAYA, O. A.; PAUKOV, I. E.; SOKOLOVA, S. I. Thermodynamic properties of gallium from 5 to 320 K . 431

GOVORCHIN, S. W.; KANA'AN, A. S. Sublimation study of triphenyl aluminum 437

CHRISTENSEN, J. J.; WALKER, T. A. C.; SCHOFIELD, R. S.; FAUX, P. W.; HARDING, P. R.; IZATT, R. M. The excess enthalpies of (carbon dioxide + hexane) at 308.15, 358.15, and 413.15 K from 7.50 to 12.50 MPa . . 445

CHYLINSKI, K.; STRYJEK, R. Excess volumes of quinoline + *n*-hexane, + *n*-octane, + *n*-decane, + *n*-undecane, and + *n*-tetradecane at 298.15 K. 453

AMBROSE, D.; PATEL, N. C. The correlation and estimation of vapour pressures. IV. Extrapolation of vapour pressures and estimation of critical pressures by the principle of corresponding states using two reference fluids with non-spherical molecules 459

FERNÁNDEZ-FASSNACHT, E.; RÍO, F. DEL. The vapour pressure of CO_2 from 194 to 243 K 469

UCHYTILOVÁ, V.; MAJER, V.; SVOBODA, V.; HEMER, I. Enthalpies of vaporization and cohesive energies of 2-chloro-1,1,2-trifluoroethyl ethyl ether, 2-chloro-1,1,2-trifluoroethyl propyl ether, 2-chloro-1,1,2-trifluoroethyl chloromethyl ether, and 2-chloro-1,1,2-trifluoroethyl difluoromethyl ether 475

CLARK, G. M.; NELSON, P. G.; MARTIN, J. F. Heat capacity of potassium hexafluoroferrate(III) between 12 and 320 K 481

GATHER, B.; BLACHNIK, R. A simple heat-flow calorimeter and the excess enthalpies of (gold + gallium) and (gold + indium). 487

CRESPO COLIN, A.; COMPOSTIZO, A.; DÍAZ PEÑA, M. Excess Gibbs energy and excess volume of (cyclohexane + 2-propanone) and of (cyclohexane + 2-butanone) 497

Number 6, June 1984

CHHOR, K.; POMMIER, C.; DIOT, M. Low-temperature thermodynamic study of the stable and metastable phases of $(C_4H_4N)Fe(C_5H_5)$, $(C_4H_4P)Fe(C_5H_5)$, and $(C_5H_5)_2Cr$ crystals 503

LAVUT, E. G.; TIMOFEYEV, B. I.; YULDASHEVA, V. M.; GALCHENKO, G. L. Enthalpy of formation of nickel chloride 519

MEISINGSET, K. K.; GRØNVOLD, F. Thermodynamic properties and phase transitions of salt hydrates between 270 and 400 K. III. $CH_3CO_2Na \cdot 3H_2O$, $CH_3CO_2Li \cdot 2H_2O$, and $(CH_3CO_2)_2Mg \cdot 4H_2O$. . 523

HAMAM, S. E. M.; KUMARAN, M. K.; BENSON, G. C. Excess enthalpies and excess volumes of each of the mixtures: (*n*-dodecane + an isomer of hexane) at 298.15 K 537

SMITH, G. R.; WORMALD, C. J. The excess molar enthalpies of $\{xH_2O+(1-x)CO\}(g)$ and $\{xH_2O+(1-x)CO_2\}(g)$ 543

GREGORCZYK, Z.; JURZYK, E. Thermodynamic properties of liquid (indium + bismuth + lead) by a potentiometric method 551

AZPIAZU, Y.; ROYO, F.; GUTIÉRREZ LOSA, C. Vapour pressures of (tetrachloromethane + α,ω-dichloroalkane) 561

MAJER, V.; SVOBODA, V.; PECHÁČEK, J.; HÁLA, S. Enthalpies of vaporization and cohesive energies of eight C_9 to C_{11} alkanes 567

INABA, H.; TAKAHASHI, S.; MIMA, T.; NAITO, K. A high-temperature Tian-Calvet type calorimeter and an analysis of the baseline fluctuation . . 573

JASRA, R. V.; AHLUWALIA, J. C. Enthalpies and heat capacities of dissolution, apparent molar heat capacities, and apparent molar volumes of some mono, di-, tri-, and tetra-saccharides in water 583

TAKAGI, T.; TERANISHI, H. Ultrasonic speed in (chlorobenzene + nitrobenzene) under high pressures 591

NOTES

REDDY, K. S. Excess volumes of (benzene + 2-ethoxyethanol) at 303.15, 313.15, and 323.15 K. 597

KUMARAN, M. K.; BENSON, G. C. Excess molar volume of (benzene + *n*-hexane) at 298.15 and 323.15 K 599

SCHOTSCH, K.; WOLF, B. A. Excess volumes of (*t*-butyl chloride + tetrachloromethane) and of (cyclohexane + 1,3-diphenylbutane) . . 601

Number 7, July 1984

GARCIA, M.; REY, C.; PEREZ VILLAR, V.; RODRIGUEZ, J. R. Excess volumes of (*n*-heptane + *n*-octane + cyclohexane) at 298.15 K 603

KALINOWSKA, B.; WÓYCICKI, W. Heat capacities of liquids in the temperature interval between 90 and 300 K and at atmospheric pressure. III. Heat capacities and excess heat capacities of (*n*-hexan-1-ol + *n*-hexane) . . 609

OTSUKA, S.; HAMA, N.; KOZUKA, Z. Activity of oxygen in liquid (lead + thallium) alloys 615

HANDA, Y. P.; HAWKINS, R. E.; MURRAY, J. J. Calibration and testing of a Tian-Calvet heat-flow calorimeter. Enthalpies of fusion and heat capacities for ice and tetrahydrofuran hydrate in the range 85 to 270 K . 623

KIRKLIN, D. R.; DOMALSKI, E. S. Enthalpy of combustion of purine . . 633

INABA, H.; MIYAHARA, K.; NAITO, K. Measurement of the molar heat capacities of MoO_2 and MoO_3 from 350 to 950 K 643

LOBO, L. Q.; STAVELEY, L. A. K. Thermodynamics of liquid (hydrogen chloride + dinitrogen oxide) 653

BICKERTON, J.; PIEDADE, M. E. M. DA; PILCHER, G. Enthalpy of formation of tetrabromomethane by rotating-bomb calorimetry 661

EWING, M. B.; KIMPTON, S. K.; MCGLASHAN, M. L. Limited azeotropy in fluid (benzene + hexafluorobenzene). 669

RAU, H.; SCHNEDLER, E. Standard molar enthalpy of formation of GeO(g) from flow measurements 673

AZEVEDO, F. G. DE; OLIVEIRA, W. A. DE. Ebulliometric behaviour of aqueous solutions of urea 683

KUMARAN, M. K.; KIMURA, F.; HALPIN, C. J.; BENSON, G. C. Excess volumes and ultrasonic speeds for (3,6-dioxaoctane + *n*-heptane) 687

SIPOWSKA, J. T.; WIECZOREK, S. A. Vapour pressures and excess Gibbs free energies of (cyclohexanol + *n*-heptane) between 303.147 and 373.278 K . 693

NOTE

GOVINDAN, A. P.; VARMA, Y. B. G.; ANANTH, M. S. Isothermal (vapour + liquid) equilibria of (tetrahydrofuran + acetonitrile + 1,2-dichloroethane) 701

Number 8, August 1984

GOVORCHIN, S. W.; KANA'AN, A. S.; KANAMUELLER, J. M. Sublimation study of triphenyl boron and the bond-dissociation enthalpy of $B–C_6H_5$ . . 703

VALERO, J.; BUENO, M. C.; PEREZ, P.; GRACIA, M.; GUTIÉRREZ LOSA, C. Excess molar volumes of several mixtures containing 1,2-dichloromethane 711

AMITIN, E. B.; NABUTOVSKAYA, O. A.; PAUKOV, I. E.; SUKHOVEY, K. S. Heat capacity of $NH_4Cl_{1-x}Br_x$ solid solutions in the vicinity of the γ-to-β phase transition 719

AWWAD, A. M.; KANBOUR, F. I.; ALLOS, E. I. Excess volumes of (an *n*-alkanol + *N*-formylmorpholine) 733

NAGATA, I. Liquid–liquid equilibria of (methanol + cyclohexane + chlorobenzene) and of (methanol + methylcyclohexane) + benzene, + ethanol, and + toluene at 298.15 K 737

DOWNIE, D. B.; MARTIN, J. F. Heat capacities of transition-metal alloys. 2. Alpha- and sigma-phase $Fe_{0.525}Cr_{0.475}$ and $Fe_{0.566}Cr_{0.434}$ . . . 743

O'HARE, P. A. G.; MALM, J. G. Thermochemistry of uranium compounds. XIV. Standard molar enthalpy of formation at 298.15 K of $AgUF_6$ and its relation to the thermodynamics of fluorination of AgF to AgF_2 . . 753

MAJER, V.; UCHYTILOVÁ, V.; SVOBODA, V.; POŠTA, A. Enthalpies of vaporization and cohesive energies of 2-chloro-1,1,2-trifluoroethyl methyl ether, 2-chloro-1,1,2-trifluoroethyl isopropyl ether, 2-chloro-1,1,2-trifluoroethyl butyl ether, and 2-chloro-1,1,2-trifluoroethyl chlorofluoromethyl ether 761

MUCKLEJOHN, S. A.; O'BRIEN, N. W. The vapour pressure and evaporation thermodynamics of sodium iodide 767

NIEPMANN, R. Thermodynamic properties of acetonitrile. 1. Speed of sound between 240 and 475 K and up to 60 MPa 779

BANNISTER, M. J. The standard molar Gibbs free energy of formation of PbO. Oxygen concentration-cell measurements at low temperatures . . . 787

Miller, R. C.; Williamson, A. G. Excess molar enthalpies for (*n*-hexane + *n*-hexadecane) and for three- and four-component alkane mixtures simulating this binary mixture 793

Jambon, C.; Merlin, J. C. Excess volumes of tributylphosphite with branched and linear alkanes 801

Number 9, September 1984

Letcher, T. M. Excess volumes of (benzene + a cycloalkane) and of (benzene + an *n*-alkane) at two temperatures 805

Wallis, K. P.; Clancy, P.; Zollweg, J. A.; Streett, W. B. Excess thermodynamic properties for $\{xCO_2+(1-x)C_2H_6\}$(l): experiment and theory 811

Smith, G. R.; Fahy, M. J.; Wormald, C. J. The excess molar enthalpy of $\{xH_2O+(1-x)C_nH_{2n+2}\}$(g) for $n = 5$, 6, 7, and 8 825

Gregorczyk, Z.; Góral, J. Thermodynamic properties of liquid (lead + silver + bismuth) by a potentiometric method 833

Węgrzyn, T. F.; Hedwig, G. R. Excess enthalpies of aqueous solutions of (urea + one of four dipeptides). 843

Niepmann, R. Thermodynamic properties of propane and *n*-butane. 2. Speeds of sound in the liquid up to 60 MPa. 851

Cobos, J. C.; Villamañan, M. A.; Casanova, C. Excess enthalpies of (*n*-alkanol + 2,5-dioxahexane) at 298.15 K 861

Mraw, S. C.; Kleppa, O. J. Calvet-type calorimeter for the study of high-temperature processes. I. Description of an apparatus appropriate for organic materials 865

Mraw, S. C.; Keweshan, C. F. Calvet-type calorimeter for the study of high-temperature processes. II. New ballistic method for the enthalpy of vaporization of organic materials at high temperatures 873

White, M. A. Phase transitions in $Mg(ClO_4)_2 \cdot 6H_2O$, as determined from the measurement of its heat capacity from 25 to 355 K by adiabatic calorimetry 885

Apelblat, A.; Tamir, A.; Wagner, M. Excess Gibbs energy of (formic acid + benzyl alcohol) and (acetic acid + benzyl alcohol) 891

Hatem, G.; Gaune-Escard, M. The unsymmetrical reciprocal molten-salt mixture $(Na^+ + Rb^+ + F^- + SO_4^{2-})$. Calorimetric measurement and evaluation of the excess molar enthalpies 897

Number 10, October 1984

CHRISTENSEN, J. J.; ZEBOLSKY, D. M.; SCHOFIELD, R. S.; CORDRAY, D. R.; IZATT, R. M. The excess enthalpies of (ethane + chlorodifluoromethane) from 293.15 to 383.15 K at 5.15 MPa 905

AWWAD, A. M.; KANBOUR, F. I.; ALLOS, E. I. Effect of molecular shape on the excess volumes of mixtures of (an isomeric octane + *n*-octane) . . 917

RARD, J. A. Solubility of $Eu(NO_3)_3 \cdot 6H_2O$ in water at 298.15 K 921

DOHNAL, V.; PATTERSON, D. Excess molar enthalpies and excess molar volumes of $xCFCl_2CF_2Cl + (1-x)CCl_4$, $+(1-x)CHCl_2CH_3$, $+(1-x)CH_2ClCH_2Cl$, $+(1-x)CCl_3CH_3$, and $+(1-x)CHCl_2CHCl_2$ at 298.15 K 927

APELBLAT, A.; WEINTRAUB, R.; TAMIR, A. Enthalpy of solution of lithium chloride and of lithium chloride monohydrate in water 935

ZHENG, G.-K.; SCHMITTER, B. M.; KNOBLER, C. M.; SCOTT, R. L. Isomer effects in mixtures of hydrocarbons. II. Some re-measurements of excess volumes 943

IMPERATORI, P.; PIACENTE, V. Vapour pressure of condensed HgSe from torsion-effusion measurements 949

NAGATA, I. Isothermal (vapor + liquid) equilibria of chlorobenzene + acetonitrile and + methanol, and of (methanol + acetonitrile + chlorobenzene) 955

AMITIN, E. B.; BESSERGENEV, W. G.; KOVALEVSKAYA, YU. A. Thermal expansion of dysprosium from 60 to 300 K 959

CHILINGAROV, N. S.; KOROBOV, M. V.; SIDOROV, L. N.; MIT'KIN, V. N.; SHIPACHEV, V. A.; ZEMSKOV, S. V. Electron affinity of rhodium tetrafluoride 965

NAGATA, I.; TAMURA, K. Excess molar enthalpies of (ethanol + tetrachloromethane) and (ethyl ethanoate + cyclohexane), and of (ethanol + benzene + tetrachloromethane) and (ethanol + ethyl ethanoate + cyclohexane) at 298.15 K 975

EVERETT, D. H.; LANGDON, A. G.; MAHER, P. Developments in immersion calorimetry: design and testing of an improved seal-breaking technique . 981

TOPOR, L.; KLEPPA, O. J. Standard molar enthalpy of formation of LaB_6 by high-temperature calorimetry 993

Number 11, November 1984

FERNÁNDEZ-FASSNACHT, E.; RÍO, F. DEL. The vapour pressure of CCl_3F from 218 to 291 K 1003

DOHNAL, V.; PATTERSON, D. Excess volumes of (cyclopentane + an isomeric hexane or heptane) at 298.15 K 1007

HAMAM, S. E. M.; KUMARAN, M. K.; BENSON, G. C. Excess enthalpies of some binary mixtures: (a C_5-alkanol + *n*-heptane) at 298.15 K . . 1013

MAJER, V.; SVOBODA, V.; LENCKA, M. Enthalpies of vaporization and cohesive energies of pyridine and isomeric methylpyridines 1019

BEYER, R. P.; JOHNSON, E. A. Heat capacity of aluminum silicon carbide (Al_4SiC_4) from 5.26 to 1047 K 1025

TAKAGI, T.; TERANISHI, H. Ultrasonic speeds and thermodynamics for (chlorobenzene + bromobenzene) under high pressures 1031

APELBLAT, A. Enthalpy of solution of rubidium sulphate in water. . . 1037

OLOFSSON, G.; OSHODJ, A. A.; QVARNSTRÖM, E.; WADSÖ, I. Calorimetric measurements on slightly soluble gases in water. Enthalpies of solution of helium, neon, argon, krypton, xenon, methane, ethane, propane, *n*-butane, and oxygen at 288.15, 298.15, and 308.15 K 1041

SOMOZA, J. A.; GARCIA, M.; REY, C.; RODRIGUEZ, J. R.; PEREZ VILLAR, V. Excess volumes of (*n*-heptane + *n*-nonane + cyclohexane) and (*n*-heptane + *n*-decane + cyclohexane) at 298.15 K 1053

MAJER, V.; SVOBODA, V.; HYNEK, V. On the enthalpy of vaporization of isomeric butanols 1059

DRAEGER, J. A. Chemical-thermodynamic properties of molecules that undergo inversion. II. The methylanilines 1067

DRAEGER, J. A. Chemical-thermodynamic properties of molecules that undergo inversion. III. The aminopyridines 1075

NAUMOWICZ, E.; WÓYCICKI, W. Excess enthalpies of gaseous mixtures containing ammonia. 1. (Ammonia + argon) 1081

OETTING, F. L.; HODGES, A. E., III; HASCHKE, J. M.; FLOTOW, H. E. The heat capacity (8 to 350 K) of $PuH_{1.9}$, heat capacity (340 to 600 K) of $PuH_{2.0}$, and recommended thermodynamic properties of PuH_2 to 600 K 1089

Number 12, December 1984

RÓG, G.; BORCHARDT, G. Thermodynamics of nickel orthosilicate . . 1103

MARSH, K. N.; ALLAN, W. A.; RICHARDS, A. E. Excess enthalpies and excess volumes of 1-nitropropane +, and 2-nitropropane +, each of several non-polar liquids 1107

COLOMINA, M.; JIMENEZ, P.; ROUX, M. V.; TURRION, C. Thermochemical properties of benzoic acid derivatives. XI. Vapour pressures and enthalpies of sublimation and formation of the six dimethylbenzoic acids 1121

CONNOLLY, J. F.; KANDALIC, G. A. Thermodynamic properties of dilute solutions of carbon monoxide in a hydrocarbon 1129

KENYON-BLAIR, E.; MORCOM, K. W. Excess enthalpy of (tetrachloromethane + 2-propanone): deuterium isotope effect 1141

COMERT, H.; PRATT, J. N. The standard molar Gibbs free energy of formation of NiO from high-temperature e.m.f. measurements . . 1145

RIBEIRO DA SILVA, M. D. M. C.; RIBEIRO DA SILVA, M. A. V.; PILCHER, G. Enthalpies of combustion of 1,2-dihydroxybenzene and of six alkylsubstituted 1,2-dihydroxybenzenes 1149

KRATZKE, H.; MÜLLER, S. Thermodynamic quantities for propane. 3. The thermodynamic behaviour of saturated and compressed liquid propane. 1157

SETTLE, J. L.; O'HARE, P. A. G. Thermochemistry of inorganic sulfur compounds. III. The standard molar enthalpy of formation at 298.15 K of $US_{1.992}$ (β-uranium disulfide) by fluorine bomb calorimetry . . 1175

KUMARAN, M. K.; BENSON, G. C.; D'ARCY, P. J.; HALPIN, C. J. Speed of sound, molar volume, and molar isobaric heat capacity for binary liquid mixtures: analysis in terms of van der Waals's one-fluid theory . . 1181

CRISTOL, B.; HOURIEZ, J.; BALESDENT, D. Standard molar enthalpies of formation of $KCdCl_3$ and K_4CdCl_6 by high-temperature drop calorimetry 1191

SANAHUJA, A.; CESARI, E. Enthalpy of solution of KCl and NaCl in water at 298.15 K 1195

INDEX TO VOLUME 16 1203